

P. Sladkov

Solitonic model of electron, proton and neutron

## Abstract

In paper, which is submitted, electron, proton and neutron are considered as spherical areas, inside which monochromatic electromagnetic wave of corresponding frequency spread along parallels, at that along each parallel exactly half of wave length for electron and proton and exactly one wave length for neutron is kept within, thus this is rotating soliton. This is caused by presence of dispersion and anisotropy of strictly defined type inside the particles. Electric field have only radial component, and magnetic field - only meridional component. By solution of corresponding edge task, functions of distribution of electromagnetic field inside the particles and on their boundary surfaces were obtained. Integration of distribution functions of electromagnetic field through volume of the particles lead to system of algebraic equations, solution of which give all basic parameters of particles: charge, rest energy, mass, radius, magnetic moment and spin.

### 1. Rotating monochromatic electromagnetic wave.

Let us write down Maxwell's equations in spherical coordinates supposing that:

- 1) there are no losses;
- 2) only  $\dot{E}_r, \dot{H}_\theta, \dot{j}_\varphi, \dot{\rho}$  are not equal to zero.

$$\frac{1}{r} \left( \frac{\partial}{\partial r} (r \dot{H}_\theta) \right) = \dot{j}_\varphi; \quad (1)$$

$$\frac{1}{r \sin \theta} \frac{\partial \dot{E}_r}{\partial \varphi} = -i \omega \mu \dot{H}_\theta; \quad (2)$$


$$\frac{1}{r} \frac{\partial \dot{E}_r}{\partial \theta} = i \omega \mu \dot{H}_\varphi = 0; \quad (3)$$

$$\frac{1}{r \sin \theta} \left( -\frac{\partial \dot{H}_\theta}{\partial \varphi} \right) = i \omega \varepsilon \dot{E}_r; \quad (4)$$

$$\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \varepsilon \dot{E}_r) = \dot{\rho}; \quad (5)$$

$$\frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta (\mu \dot{H}_\theta)) = 0. \quad (6)$$

Here  $r, \theta, \varphi$  - spherical coordinates of the observation point;  $\dot{E}_r$  и  $\dot{H}_\theta$  - components of the electromagnetic field,  $\dot{j}_\varphi$  - density of electric current,  $\dot{\rho}$  - volume charge density;  $\omega$  - circular frequency of field alteration,  $i$  - imaginary unit,  $\varepsilon$  - dielectric permittivity,  $\mu$  - magnetic permeability.



**Fig. 1**

Substituting the expression for  $\dot{H}_\theta$  from (2) in (4), we obtain:

$$\frac{\partial^2 \dot{E}_r}{\partial \varphi^2} + \varepsilon \mu \omega^2 r^2 \sin^2 \theta \dot{E}_r = 0; \quad (7)$$

$$\frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \dot{E}_r}{\partial \varphi^2} + \omega^2 \varepsilon \mu \dot{E}_r = 0; \quad (7')$$

This is Helmholtz homogeneous equation. Let us designate

$$\sqrt{\varepsilon \mu} \omega r \sin \theta = k_1 - \quad (7'')$$

wave number. General solution of Helmholtz equation:

$$\dot{E}_r = E_0 e^{-ik_1 \varphi} + E_0 e^{ik_1 \varphi}. \quad (8)$$

This expression describes two waves, moving to meet one another by circular trajectories, along the parallels. Pointing's vector in each point is directed at tangent to the corresponding parallel.

Let us consider a wave, moving in positive direction  $\varphi$ .

$$\dot{E}_r = E_0 e^{-ik_1 \varphi} F(r, \theta); \quad (9)$$

Here

$$k_1 \varphi = \sqrt{\varepsilon \mu} (\omega r \sin \theta) \varphi -$$

wave phase;

$k_1$  – dimensionless analog of the wave number. If to introduce a wave number of traditional dimension ( $\frac{1}{m}$ );

$$\beta = \omega \sqrt{\varepsilon \mu} = \frac{k_1}{r \sin \theta},$$

the wave phase will be written down as

$$k_1 \varphi = \sqrt{\varepsilon \mu} \omega (r \sin \theta) \varphi = \beta l,$$

where

$$l = (r \sin \theta) \varphi -$$

arc length along the corresponding parallel. In the considered case the wave number is a function of coordinates and frequency. Thus, the wave, which is described, can exist only at availability of spatial and frequency dispersion. Dispersion equations will be obtained below, apart from the already found expression (7'').

From expression (2), taking into account (7'') and (9), we have:

$$\dot{H}_\theta = \frac{\sqrt{\varepsilon \mu} \omega r \sin \theta}{\mu \omega r \sin \theta} E_0 e^{-ik_1 \varphi} F(r, \theta) = \frac{E_0}{z} e^{-ik_1 \varphi} F(r, \theta) . \quad (9')$$

For actual amplitudes:

$$E_r = E_0 F(r, \theta) \sin k_1 \varphi ; \quad (10)$$

$$H_\theta = \frac{E_0}{z} F(r, \theta) \sin k_1 \varphi . . \quad (10')$$

Here

$$z = \sqrt{\frac{\mu}{\varepsilon}} -$$

means characteristic impedance.

The last expressions describe an electromagnetic wave, rotating around axis Z in positive direction  $\varphi$ . Conditions of self-consistency:

- 1)  $z = const$ ;
- 2) along each parallel on the circle length, the integer number of half-waves must be kept within.

$$2\pi r \sin \theta = n \frac{\lambda}{2} ; \quad (11)$$

here  $\lambda = \frac{v}{f}$  -

wave length,  $v$  - phase velocity of wave,  $f$  - frequency,  $n = 1, 2, 3, \dots$

Let us consider the case when  $n = 1$ ,

$$2\pi r \sin \theta = \frac{\lambda}{2} ;$$

$$v = 2\omega r \sin \theta. \quad (11')$$

Along each parallel, exactly half of wave length is kept within.

Phase velocity of wave is the function of frequency and distance up to the axis of rotation.

$$v = \frac{1}{\sqrt{\varepsilon \mu}} = 2\omega r \sin \theta ;$$

$$\varepsilon \mu = \frac{1}{4\omega^2 r^2 \sin^2 \theta} ; \quad (11'')$$

$$z = \sqrt{\frac{\mu}{\varepsilon}} ;$$

$$\mu = \varepsilon z^2; \quad (11''')$$

we are substituting in (11''):

$$z = \frac{1}{2\omega r \varepsilon \sin \theta}; \quad (12)$$

$$\varepsilon = \frac{1}{2\omega r z \sin \theta}. \quad (12')$$

From (11'')  $\varepsilon = \frac{\mu}{z^2}$ ;

we are substituting in (12').

$$\mu = \frac{z}{2\omega r \sin \theta}; \quad (12'')$$

$$z = 2\mu \omega r \sin \theta. \quad (12''')$$

Taking into account (8) and (11'')

$$k_1 = \frac{\omega r \sin \theta}{2\omega r \sin \theta} = \frac{1}{2}.$$

Then

$$E_r = E_0 F(r, \theta) \sin \frac{\varphi}{2}; \quad (13)$$

$$H_\theta = \frac{E_0}{z} F(r, \theta) \sin \frac{\varphi}{2}. \quad (13')$$

Function  $\sin \frac{\varphi}{2}$  is onevalued in angles interval  $0 \leq \varphi \leq 2\pi$ .

This situation can be interpreted as rotation of spherical coordinate system around axis z in positive direction  $\varphi$  with angular velocity  $\frac{d\varphi}{dt}$ . Let us find it from the condition

$$\omega t - \frac{\varphi}{2} = const.$$

Having differentiated this expression on t, we receive

$$\frac{d\varphi}{dt} = 2\omega.$$

At the same time the electromagnetic field, about spherical coordinate system, is determined by expressions (13) и (13').

Further from (3) : as  $\dot{H}_\varphi = 0$ ,

$$\dot{E}_r(\theta) = const;$$

$$E_r(\theta) = \text{const} \quad . \quad (14)$$

From equation (6)

$$\frac{\partial}{\partial \theta} \left( \frac{\sin \theta (z H_\theta)}{2 \omega r \sin \theta} \right) = 0$$

follows

$$\dot{H}_\theta(\theta) = \text{const} ;$$

$$H_\theta(\theta) = \text{const} . \quad (14')$$

To receive field dependence from  $r : E_r(r) ; H_\theta(r)$ , let us find solution of three-dimensional Helmholtz equation in spherical coordinates.

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial E_r}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial E_r}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 E_r}{\partial \varphi^2} + k^2 E_r = 0. \quad (15)$$

$E_r$  does not depend on  $\theta$ , look (14), therefore three-dimensional Helmholtz equation transfers into two-dimensional one.

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial E_r}{\partial r} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 E_r}{\partial \varphi^2} + k^2 E_r = 0. \quad (15')$$

Let us suppose that

$$k^2 = k_2^2 + k_3^2 ,$$

now

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial E_r}{\partial r} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 E_r}{\partial \varphi^2} + k_2^2 E_r + k_3^2 E_r = 0. \quad (15'')$$

This equation can be satisfied, if

$$\left\{ \begin{array}{l} \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 E_r}{\partial \varphi^2} + k_2^2 E_r = 0 ; \\ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial E_r}{\partial r} \right) + k_3^2 E_r = 0 . \end{array} \right. \quad (16),(17)$$

Thus, initial Helmholtz equation has split into the system of two equations. We substitute in these equations instead of  $E_r(r, \varphi) = f(r) g(\varphi)$ ,

(i.e. we are searching the solution as the product of two functions) and divide the first equation by  $f(r)$ , and the second - by  $g(\varphi)$ . We receive

$$\left. \begin{aligned} \frac{d^2 g}{d \varphi^2} + k_2^2 r^2 (\sin^2 \theta) g &= 0; \\ r^2 \frac{d^2 f}{d r^2} + 2r \frac{d f}{d r} + k_3^2 r^2 f &= 0. \end{aligned} \right\} \quad (18), (19)$$

Equations (16) and (18) are equivalent to equations (7) и (7'), which were received earlier from Maxwell's equations, and

$$k_2 = \omega \sqrt{\varepsilon \mu} = \frac{\omega}{v} = \frac{1}{2r \sin \theta};$$

$$k_1 = k_2 r \sin \theta = \frac{1}{2}$$

The solution of equation (18) was found earlier, look (13).

$$g(\varphi) = \sin \frac{\varphi}{2}. \quad (20)$$

Let us copy (19) as :

$$r^2 \frac{d^2 f}{d r^2} + 2r \frac{d f}{d r} + k_4^2 f = 0; \quad (19')$$

where

$$k_4 = k_3 r.$$

(19') – centrally symmetric Helmholtz equation. Let us suggest,

$$k_3 = \frac{\omega}{v_r},$$

where  $v_r$  – phase velocity of electromagnetic wave in radial direction. As in the central symmetric equation angular dependence is absent, it is logical to assume that

$$v_r = v = 2\omega r \sin \theta$$

at  $\theta = \frac{\pi}{2}$ ; i.e.

$$v_r = 2\omega r;$$

$$k_3 = \frac{\omega}{v_r} = \frac{1}{2r}; \quad (21)$$

$$k_4 = k_3 r = \frac{1}{2}. \quad (21')$$

Instead of (19'), we are having

$$r^2 \frac{d^2 f}{d r^2} + 2r \frac{d f}{d r} + \frac{1}{4} f = 0. \quad (19'')$$

This is Euler equation, it has the solution

$$f = r^{-\frac{1}{2}}(C_1 + C_2 \ln r). \quad (22)$$

Let us converse expression (22).

$$f = \frac{(C_1 + C_2 \ln r)}{\sqrt{r}} = \sqrt{\frac{a}{r}}(C_3 + C \ln r) = \sqrt{\frac{a}{r}}(1 + \ln C_5 + \ln r^C) = \sqrt{\frac{a}{r}}(1 + \ln C_5 r^C). \quad (22')$$

Here  $C_1 = \sqrt{a} C_3$ ;  $C_2 = \sqrt{a} C$ ;  $C_3 = 1 + \ln C_5$ ;  $a$  - value of radius  $r$ , at which the rotating monochromatic electromagnetic wave ceases to exist, and  $E_r = E_0$ ;  $f = 1$ ; hence

$$\begin{aligned} 1 + \ln C_5 a^C &= 1; \\ \ln C_5 a^C &= 0; \\ C_5 a^C &= 1; \\ C_5 &= \frac{1}{a^C}. \end{aligned} \quad (22'')$$

In view of this,

$$f = \sqrt{\frac{a}{r}} (1 + \ln (\frac{r}{a})^C) = \sqrt{\frac{a}{r}} (1 + C \ln \frac{r}{a}).$$

Let us designate  $C = p$ ; now

$$f = \sqrt{\frac{a}{r}} (1 + p \ln \frac{r}{a}).$$

Thus, for  $E_r$  we are having

$$E_r = E_0 g(\varphi) f(r) = E_0 \sqrt{\frac{a}{r}} (1 + p \ln \frac{r}{a}) \sin \frac{\varphi}{2}. \quad (23)$$

At  $r \rightarrow \infty$ ,  $E_r = 0$ ;  $f = 0$ .

Really

$$\lim_{r \rightarrow \infty} \frac{\ln r}{\sqrt{r}} = \lim_{r \rightarrow \infty} \frac{1/r}{1/2 \sqrt{r}} = 0.$$

So that at alteration of  $r$  within the interval from 0 to  $a$ ,  $E_r$  would not change its sign, observance of the following requirement is necessary:  $p \leq 0$ .

At  $r = 0$ ,  $E_r = \infty$ ;  $f = \infty$ .

At  $r = a$ ,  $f = 1$ ;  $E_r = E_0$ .

## 2. System of equations for electron.

Basing on results of the previous section, let us write down expressions for electromagnetic field inside the electron, assuming that it is concentrated inside the orb of radius  $a$ .

$$E_r = E_0 \sqrt{\frac{a}{r}} \left(1 + p \ln \frac{r}{a}\right) \sin \frac{\varphi}{2}; \quad (23')$$

$$H_\theta = \frac{E_0}{z} \sqrt{\frac{a}{r}} \left(1 + p \ln \frac{r}{a}\right) \sin \frac{\varphi}{2}. \quad (23'')$$

Here  $a$  is electron radius,  $E_0$  - amplitude of electric field intensity at  $r = a$ ;  $z = const$  - characteristic impedance inside the electron,  $p$  - unknown coefficient and  $p \leq 0$ .

At that the internal electron medium possesses frequent and spatial dispersion, as well as anisotropy. Dispersion equations have the following appearance.

$$v_r = v_\theta = 2 \omega r; \quad (24)$$

$$v_\varphi = 2 \omega r \sin \theta; \quad (24')$$

$$z_r = z_\theta = z_\varphi = z = const. \quad (24'')$$

Here  $v_r, v_\theta, v_\varphi$  - phase velocity of rotating monochromatic electromagnetic wave in corresponding direction. In viewed case, the electromagnetic wave is being spread only in the direction of  $\varphi$ , and we shall need expressions  $v_r$  and  $v_\theta$  for searching the formulas of dielectric and magnetic permeability, as well as wave numbers of corresponding directions;  $z_r, z_\theta, z_\varphi, z$  - characteristic impedances inside the electron;  $\varepsilon_\varphi$  и  $\mu_\varphi$  were found before, see (12'), (12'').

$$\varepsilon_\varphi = \frac{1}{2 \omega r z \sin \theta} = \frac{1}{v_\varphi z};$$

$$\mu_\varphi = \frac{z}{2 \omega r \sin \theta} = \frac{z}{v_\varphi}.$$

In view of (24), (24'), (24''), let us write down expressions for  $\varepsilon_r, \varepsilon_\theta, \mu_r, \mu_\theta$ .

$$\varepsilon_r = \frac{1}{2 \omega r z};$$

$$\mu_r = \frac{z}{2 \omega r};$$

$$\varepsilon_\theta = \frac{1}{2 \omega r z};$$

$$\mu_\theta = \frac{z}{2 \omega r}. \quad (24''')$$

From considerations and formulas adduced, it follows that dielectric and magnetic permeability are tensor values.

$$\|\varepsilon\| = \begin{vmatrix} \varepsilon_r & 0 & 0 \\ 0 & \varepsilon_\varphi & 0 \\ 0 & 0 & \varepsilon_\theta \end{vmatrix} = \begin{vmatrix} \frac{1}{2 \omega r z} & 0 & 0 \\ 0 & \frac{1}{2 \omega r z \sin \theta} & 0 \\ 0 & 0 & \frac{1}{2 \omega r z} \end{vmatrix}.$$

$$\|\mu\| = \begin{vmatrix} \mu_r & 0 & 0 \\ 0 & \mu_\varphi & 0 \\ 0 & 0 & \mu_\theta \end{vmatrix} = \begin{vmatrix} \frac{z}{2 \omega r} & 0 & 0 \\ 0 & \frac{z}{2 \omega r \sin \theta} & 0 \\ 0 & 0 & \frac{z}{2 \omega r} \end{vmatrix}.$$

Let us find dimensionless wave numbers.

$$k_\varphi = \frac{\omega}{v_\varphi} r \sin \theta = \frac{\omega r \sin \theta}{2 \omega r \sin \theta} = \frac{1}{2};$$

$$k_\theta = \frac{\omega}{v_\theta} r = \frac{\omega r}{2 \omega r} = \frac{1}{2};$$

$$k_r = \frac{\omega}{v_r} r = \frac{\omega r}{2 \omega r} = \frac{1}{2}.$$

Thus

$$k_\varphi = k_\theta = k_r = \frac{1}{2}.$$

Let us remind that in the viewed case, the electromagnetic wave is spread only in the direction of  $\varphi$ .

At  $r = 0$  we are having a special point:

$$E_r = \infty; H_\theta = \infty; \rho = \infty; j_\varphi = \infty; \|\varepsilon\| = \infty; \|\mu\| = \infty.$$

Despite of this, all basic electron's parameters - charge  $q$ , rest energy  $W$ , magnetic moment  $M$ , - expressed through integrals by volume from the functions specified above, prove to be finite quantities. See further.

From (5), we find volume charge density inside electron  $\rho$ .

$$\begin{aligned} \rho = \operatorname{div}(\varepsilon_r E_r) &= \frac{1}{r^2} \frac{\partial}{\partial r} \left[ \frac{r^2 E_0 \sin \frac{\varphi}{2}}{2 \omega r z} \left( \left( \frac{a}{r} \right)^{\frac{1}{2}} + p \left( \frac{a}{r} \right)^{\frac{1}{2}} \ln \frac{r}{a} \right) \right] = \\ &= \frac{E_0 \sqrt{a} \sin \frac{\varphi}{2}}{2 \omega z r^2} \left( \left( \frac{1}{2} + p \right) \frac{1}{\sqrt{r}} + \frac{p}{2\sqrt{r}} \ln \frac{r}{a} \right). \end{aligned} \quad (25)$$

Integrating  $\rho$  on electron's volume, we shall receive this expression for its charge  $q$ .

$$\begin{aligned} q = \int_V \rho dV &= \frac{E_0 \sqrt{a}}{2 \omega z} \int_0^{2\pi} \int_0^\pi \int_0^a \frac{\sin \frac{\varphi}{2} r^2 \sin \theta}{r^2} \left( \left( \frac{1}{2} + p \right) \frac{1}{\sqrt{r}} + \frac{p}{2\sqrt{r}} \ln \frac{r}{a} \right) d\varphi d\theta dr = \\ &= \frac{4 E_0 \sqrt{a}}{\omega z} (2\sqrt{a} \left( \frac{1}{2} + p \right) + p\sqrt{a} \ln a - 2p\sqrt{a} - p\sqrt{a} \ln a - p\sqrt{0} \ln 0) = \frac{4 E_0 a}{\omega z}. \end{aligned} \quad (26)$$

On the other hand, from the third integral Maxwell's equation, it is possible to find electron's charge as a stream of vector electric induction  $D$  through the surface of the orb of radius  $a$ .

$$q = \oint_S \varepsilon_r E_r dS = \int_0^{2\pi} \int_0^\pi \frac{E_0 \sin \frac{\varphi}{2} a^2 \sin \theta}{2 \omega z a} d\varphi d\theta = \frac{4 E_0 a}{\omega z}. \quad (26')$$

As we can see, expressions (26) и (26') are equivalent to each other.

From (1), we obtain expression for current density  $j_\varphi$ .

$$j_\varphi = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{E_0 \sin \frac{\varphi}{2}}{z} \left( \sqrt{\frac{a}{r}} + p \sqrt{\frac{a}{r}} \ln \frac{r}{a} \right) \right) = \frac{E_0 \sqrt{a} \sin \frac{\varphi}{2}}{r z} \left( \left( \frac{1}{2} + p \right) \frac{1}{\sqrt{r}} + \frac{p}{2\sqrt{r}} \ln \frac{r}{a} \right). \quad (27)$$

From expressions (25), (27) it is visible that in the interval of change of  $r$  from 0 to  $a$ ,  $\rho$  and  $j_\varphi$  once change the sign. It can be explained by the fact that in the viewed structure, the substantial role is played by the rotating monochromatic electromagnetic wave, and the space charge density and electric current density – are auxiliary or even fictitious quantities in the sense that inside the particle there is neither any charged substance nor its motion. Inside the electron, it is not the charge that is the source of electric field, but electric field is the source of the charge. In its turn, it is not the electric current that is the

source of magnetic field, but magnetic field is the source of the electric current. Thus, a deduction about vector nature of elementary charge can be made.

Now we shall determine electron's rest energy as electromagnetic wave energy inside a particle.

$$W = \int_V w dV.$$

Here  $w$  - is volume density of electromagnetic wave energy,

$$w = \frac{\Pi}{v_\varphi}, \text{ where}$$

$\Pi$  – Pointing vector,

$$\Pi = [E_r, H_\theta],$$

$v_\varphi$  - phase velocity of electromagnetic wave in direction of  $\varphi$  .

$$v_\varphi = 2 \omega r \sin \theta.$$

$$\begin{aligned} W &= \int_0^{2\pi} \int_0^\pi \int_0^a \frac{E_0^2 \sin^2 \frac{\varphi}{2}}{2 \omega r z \sin \theta} \left( \frac{a}{r} + 2p \frac{a}{r} \ln \frac{r}{a} + p^2 \frac{a}{r} \ln^2 \frac{r}{a} \right) r^2 \sin \theta d\varphi d\theta dr = \\ &= \frac{\pi^2 E_0^2 a}{2 \omega z} (a + 2 p a \ln a - 2 p 0 \ln 0 - 2 p a - 2 p a \ln a + p^2 a (2 - 0 \ln^2 0 + 2 * 0 \ln 0)) = \\ &= \frac{\pi^2 E_0^2 a^2}{2 \omega z} (1 - 2 p + 2 p^2). \end{aligned} \quad (28)$$

$$\frac{\pi^2 E_0^2 a^2}{2 \omega z} (1 - 2 p + 2 p^2) = \hbar \omega ; \quad (28')$$

here  $\hbar$  is Planck's constant.

We shall be searching electron's magnetic moment in the form of a sum.

$$M = M_m + M_L,$$

where  $M_m$  – is magnetic moment, created by volumetric current;  $M_L$  – magnetic moment, attributed to impulse moment, i.e. to rotation.

$$M_L = \gamma L,$$

where  $\gamma$  – gyromagnetic ratio;  $L$  – impulse moment of electron.

Basing on Barnett effect, we are making a supposition, that the impulse moment, attributed to rotation, creates additional magnetic moment.

Being aware of the fact that electron's impulse moment is equal  $\frac{\hbar}{2}$ , from (28') we find expression for L.

$$L = \frac{\pi^2 E_0^2 a^2}{4 \omega^2 z} (2p^2 - 2p + 1);$$

$$M_L = \frac{\gamma \pi^2 E_0^2 a^2 (2p^2 - 2p + 1)}{4 \omega^2 z};$$

or  $M_L = \gamma \frac{\hbar}{2}$ .

Let us calculate  $M_m$  as electric current magnetic moment in volume V, relating to axis z by the formula:

$$M_m = \frac{1}{2} \int_V [r_z \cdot j_\varphi] dV.$$

See for instance [3], page 111, where  $r_z$  - distance to axis z,

$$r_z = r \sin \theta.$$

$$M_m = \frac{1}{2} \int_0^{2\pi} \int_0^\pi \int_0^a \frac{r \sin \theta E_0 \sqrt{a} \sin \frac{\varphi}{2} r^2 \sin \theta}{z r^{\frac{3}{2}}} \left( p + \frac{1}{2} + \frac{p}{2} \ln \frac{r}{a} \right) d\varphi d\theta dr = \frac{\pi E_0 a^3 (8p + 5)}{z}. \quad (29)$$

$$M = M_m + M_L = \frac{\pi E_0 a^3 (8p + 5)}{z} + \gamma \frac{\hbar}{2}. \quad (29')$$

Or

$$M = \frac{\pi E_0 a^2}{z} \left[ a \left( \frac{8p + 5}{25} \right) + \frac{\gamma \pi E_0}{4 \omega^2} (1 - 2p + 2p^2) \right]. \quad (29'')$$

Thus, we have received the system of algebraic equations for electron.

$$\left. \begin{aligned} \frac{4 E_0 a}{\omega z} &= -e; & (30) \\ \frac{\pi E_0 a^3 (8p + 5)}{z} + \frac{\gamma \hbar}{2} &= -1,0011595 \frac{e \hbar}{2 m}; & (31) \\ \frac{\pi^2 E_0^2 a^2}{2 \omega z} (1 - 2p + 2p^2) &= \hbar \omega; & (32) \end{aligned} \right\}$$

Here  $e$  - charge of electron,  $m$  - its mass.

Three equations contain five unknown quantities:  $E_0, a, z, p, \gamma$ . Let us add this system with equations, which we shall receive from boundary conditions.

At  $r = a; R = a$ :

$$\varepsilon_r E_0 = \varepsilon_0 E_{\text{внешн.}} \quad (33)$$

In the exterior area, the same as and in the interior area, electric field intensity possesses only radial component. Here  $R$  - distance from electron's center to the observation point in the exterior area,  $\varepsilon_0$  - vacuum dielectric permeability.

$$\text{Further. } H_0 = \frac{E_0}{z} = H_{\text{внешн.}} \quad (34)$$

In the exterior area, the same as and in the interior area, magnetic field intensity possesses only meridional component.

It is obvious that

$$\varepsilon_r \geq \varepsilon_0, \quad (33')$$

then from (33) follows:

$$E_0 \leq E_{\text{внешн.}} \quad (33'')$$

On the other hand it is known that the electric field, having passed through dielectric layer, cannot increase, therefore

$$E_0 \geq E_{\text{внешн.}} \quad (33''')$$

In other words, correlations (33'), (33''), (33''') will be simultaneously executed only in one case, if

$$\varepsilon_r = \varepsilon_0; \quad (35)$$

$$E_{\text{внешн.}} = E_0. \quad (36)$$

Now under Biot-Savart's law, we are finding magnetic field in the exterior area.

$$B_{\text{внешн.}} = \frac{1}{4\pi} \int_V \frac{[j_\varphi R] \mu_\varphi}{R^3} dV.$$

In last expression we substitute (12'') and (27).

$$\begin{aligned} B_{\text{внешн.}} &= \frac{E_0 \sqrt{a}}{4\pi R^2 z} \int_0^{2\pi} \int_0^\pi \int_0^a \frac{z \sin \frac{\varphi}{2} \left[ \left( \frac{1}{2} + p \right) \frac{1}{\sqrt{r}} + \frac{p}{2\sqrt{r}} \ln \frac{r}{a} \right]}{2\omega r^2 \sin \theta} r^2 \sin \theta d\varphi d\theta dr = \\ &= \frac{E_0 \sqrt{a}}{2\omega R^2} \left[ \left( \frac{1}{2} + p \right) 2\sqrt{a} - p\sqrt{a} \ln a + p\sqrt{a} \ln a - p\sqrt{0} \ln 0 - 2p\sqrt{a} \right] = \frac{E_0 a}{2\omega R^2}. \end{aligned} \quad (37)$$

$$H_{\text{внешн.}} = \frac{B_{\text{внешн.}}}{\mu_0} = \frac{E_0 a}{2 \mu_0 \omega R^2}. \quad (38)$$

At  $r = a; R = a$

$$\begin{aligned} H_{\text{внутр.}} &= \frac{E_0}{z} = H_{\text{внешн.}} \\ \frac{E_0}{z} &= \frac{E_0 a}{2 \mu_0 \omega a^2} = \frac{E_0}{2 \mu_0 \omega a}; \\ z &= 2 \mu_0 \omega a. \end{aligned} \quad (39)$$

On the other hand, from (24''')

$$z = 2 \mu_\theta \omega r.$$

At  $r = a$

$$z = 2 \mu_\theta \omega a. \quad (39')$$

We substitute in (39).

$$\begin{aligned} 2 \mu_\theta \omega a &= 2 \mu_0 \omega a; \\ \mu_\theta &= \mu_0. \end{aligned} \quad (40)$$

Thus, at  $r = a$ ,

$$\begin{aligned} \varepsilon_r &= \varepsilon_\theta = \varepsilon_0; \\ \mu_\theta &= \mu_r = \mu_0; \\ v_r = v_\theta &= 2\omega a = \frac{1}{\sqrt{\mu_0 \varepsilon_0}} = c. \end{aligned} \quad (41)$$

Here  $c$  - velocity of light,  $\omega = 7,7634421 * 10^{20}$  Hz - Compton circular frequency of electron.

$$a = \frac{c}{2\omega} = 0,1930796 * 10^{-12} \text{ (m)}. \quad (42)$$

As it is known, atom's radius approximately equals to  $10^{-10}$  m, volume of atom -  $4,18879 * 10^{-30}$  m<sup>3</sup>. We found, that radius of electron equals to  $1,930796 * 10^{-13}$  m, volume of electron -  $3,0150724 * 10^{-38}$  m<sup>3</sup>. That is one electron occupies  $0,7197955 * 10^{-8}$  from atom's volume and, for example, 100 electrons (as in atoms located at the end of the periodic system) occupy  $0,7197955 * 10^{-6}$  from atom's volume.

We substitute (42) в (39).

$$z = \frac{2\mu_0 \omega c}{2\omega} = \sqrt{\frac{\mu_0}{\varepsilon_0}} = 376,73032 \text{ (Ohm)}. \quad (43)$$

Let us solve the system (30), (31), (32), taking into account (42) and (43).

$$\frac{4 E_0 c}{\omega \sqrt{\frac{\mu_0}{\varepsilon_0}} 2\omega} = -e;$$

$$E_0 = -\frac{\omega^2 \mu_0 e}{2}. \quad (30')$$

$$\frac{\pi E_0}{8\omega^3 \mu_0^2 \varepsilon_0} \left(\frac{8p+5}{25}\right) + \frac{\gamma \pi^2 E_0^2 (1-2p+2p^2)}{16\omega^4 \mu_0^{3/2} \varepsilon_0^{1/2}} = -1,0011595 \frac{e\hbar}{2m}. \quad (31')$$

$$\frac{\pi^2 E_0^2}{8\omega^3 \mu_0^{3/2} \varepsilon_0^{1/2}} (1-2p+2p^2) = \hbar\omega. \quad (32')$$

We substitute (30') in (32').

$$p^2 - p + \frac{1}{2} - \frac{16 \hbar \varepsilon_0^{1/2}}{\pi^2 e^2 \mu_0^{1/2}} = 0;$$

$$p_1 = 4,6747427;$$

$$p_2 = -3,6747427.$$

$p$  must be negative, therefore we select

$$p_2 = p = -3,6747427.$$

We substitute (30') in (31').

$$-\frac{\pi e}{16\omega \mu_0 \varepsilon_0} \left(\frac{8p+5}{25}\right) + \frac{\pi^2 \gamma e^2 \mu_0^{1/2}}{64\varepsilon_0^{1/2}} (1-2p+2p^2) = -1,0011595 \frac{e \hbar}{2m}. \quad (31'')$$

We substitute  $p$  meaning in (31'') and find  $\gamma$ .

$$\gamma = -0,2434911 * 10^{12} \left(\frac{1}{T * s}\right).$$

From solution of equation (31), it is visible that two components of magnetic moment of electron  $M_m$  и  $M_L$  are directed to opposite sides and  $M_L \succ M_m$ .

Let us also calculate numerical value of  $E_0$  by formula (30').

$$E_0 = -6,0673455 * 10^{16} \left(\frac{V}{m}\right).$$

"Dimensions" of electron for the present are not discovered by experimental way, though precision of measuring is led to  $10^{-18}$  m. Within the framework of the model considered it may be explained by the next way: electron is not hard particle with this quantity of vector E, which exist inside it, unlike from proton and neutron, quantity of vector E inside which approximately  $10^7$  times as much. Look below.

For positron, the system of equations will take a somewhat different view.

$$\left. \begin{aligned} \frac{4 E_0 a}{\omega z} &= e; & (44) \\ \frac{\pi E_0 a^3}{z} \left( \frac{8p+5}{25} \right) + \frac{\gamma \pi^2 E_0^2 a^2}{4\omega^2 z} (1-2p+2p^2) &= 1,0011595 \frac{e \hbar}{2 m}; & (45) \\ \frac{\pi^2 E_0^2 a^2}{2\omega z} (1-2p+2p^2) &= \hbar \omega; & (46) \end{aligned} \right\}$$

Boundary conditions are the same as for electron. Hence

$$z = \sqrt{\frac{\mu_0}{\varepsilon_0}};$$

$$a = \frac{c}{2\omega} = 0,1930796 * 10^{-12} (m).$$

The system of equations (44), (45), (46) with exactness to a sign, has the same solutions, as the system (30), (31), (32).

$$E_0^{e+} = -E_0^e = 6,0673455 * 10^{16} \left( \frac{V}{m} \right);$$

$$\gamma_{e+} = -\gamma_e = 0,2434911 * 10^{12} \left( \frac{1}{T * s} \right);$$

$$p_{e+} = -p_e = -3,6747427.$$

### 3. System of equations for proton.

By applying reasoning and mathematical calculations of the previous section in relation to proton, we shall receive the relevant system of equations.

$$\left. \begin{aligned} \frac{4E_0 a}{\omega z} &= e; & (47) \\ \frac{\pi E_0 a^3}{z} \left( \frac{8p+5}{25} \right) + \frac{\gamma \pi^2 E_0^2 a^2 (1-2p+2p^2)}{4\omega^2 z} &= -2,7928475 \frac{e \hbar}{2 m}; & (48) \\ \frac{\pi^2 E_0^2 a^2}{2\omega z} (1-2p+2p^2) &= \hbar \omega; & (49) \end{aligned} \right\}$$

Here corresponding letters mean parameters of proton.

Boundary conditions: at  $r = a$

$$\varepsilon_r = \varepsilon_\theta = \varepsilon_0;$$

$$\mu_\theta = \mu_r = \mu_0;$$

hence

$$z = \sqrt{\frac{\mu_0}{\varepsilon_0}};$$

$$a = \frac{c}{2\omega} = 1,0515447 * 10^{-16} \text{ m.}$$

Here  $\omega = 1,425486 * 10^{24} \text{ Hz}$  - Compton circular frequency of proton.

Solving the system (47), (48), (49), we shall receive:

$$E_0 = 2,0455794 * 10^{23} \left(\frac{\text{V}}{\text{m}}\right);$$

$$p = -3,6747427;$$

$$\gamma = -2,3081218 * 10^8 \left(\frac{1}{\text{T} * \text{s}}\right).$$

From the solution of equation (48) it is visible that two components of proton's magnetic moment  $M_m$  и  $M_L$  have identical direction, and  $M_L \succ M_m$ .

Let us write down the system of equations for antiproton.

$$\frac{4E_0 a}{\omega z} = -e; \tag{50}$$

$$\frac{\pi E_0 a^3}{z} \left(\frac{8p+5}{25}\right) + \frac{\gamma \pi^2 E_0^2 a^2 (1-2p+2p^2)}{4\omega^2 z} = 2,7928475 \frac{e \hbar}{2m}; \tag{51}$$

$$\frac{\pi^2 E_0^2 a^2}{2\omega z} (1-2p+2p^2) = \hbar\omega. \tag{52}$$

Boundary conditions: at  $r = a$

$$\varepsilon_r = \varepsilon_\theta = \varepsilon_0;$$

$$\mu_r = \mu_\theta = \mu_0;$$

hence

$$z = \sqrt{\frac{\mu_0}{\varepsilon_0}};$$

$$a = \frac{c}{2\omega} = 1,0515447 * 10^{-16} \text{ (m)}.$$

System of equations (50), (51), (52) with exactness to a sign has the same solutions, as system (47), (48), (49).

$$E_0^{\tilde{p}} = -E_0^p = -2,0455794 * 10^{23} \left(\frac{V}{m}\right);$$

$$\gamma_{\tilde{p}} = -\gamma_p = 2,3081218 * 10^8 \left(\frac{1}{T * S}\right);$$

$$p_{\tilde{p}} = p_p = p_e = -3,6747427.$$

#### 4. System of equations for neutron.

$$E_r = E_0 \sqrt{\frac{a}{r}} \left(1 + p \ln \frac{r}{a}\right) \sin \varphi; \quad (53)$$

$$H_\theta = \frac{E_0}{z} \sqrt{\frac{a}{r}} \left(1 + p \ln \frac{r}{a}\right) \sin \varphi. \quad (53')$$

Along each parallel, exactly one wave length is kept within . In this case:

$$v_\varphi = \omega r \sin \theta;$$

$$\varepsilon_\varphi = \frac{1}{\omega r z \sin \theta}; \quad (54)$$

$$\mu_\varphi = \frac{z}{\omega r \sin \theta}.$$

$$v_r = 2\omega r;$$

$$\varepsilon_r = \frac{1}{2\omega r z}; \quad (54')$$

$$\mu_r = \frac{z}{2\omega r}.$$

$$v_\theta = v_r = 2\omega r;$$

$$\varepsilon_\theta = \varepsilon_r = \frac{1}{2\omega r z}; \quad (54'')$$

$$\mu_\theta = \mu_r = \frac{z}{2\omega r}$$

In other words, anisotropy is taking place,  $\varepsilon$  and  $\mu$  are tensor quantities.

$$\|\varepsilon\| = \begin{vmatrix} \varepsilon_r & 0 & 0 \\ 0 & \varepsilon_\varphi & 0 \\ 0 & 0 & \varepsilon_\theta \end{vmatrix} = \begin{vmatrix} \frac{1}{2\omega r z} & 0 & 0 \\ 0 & \frac{1}{\omega r z \sin \theta} & 0 \\ 0 & 0 & \frac{1}{2\omega r z} \end{vmatrix}.$$

$$\|\mu\| = \begin{vmatrix} \mu_r & 0 & 0 \\ 0 & \mu_\varphi & 0 \\ 0 & 0 & \mu_\theta \end{vmatrix} = \begin{vmatrix} \frac{z}{2\omega r} & 0 & 0 \\ 0 & \frac{z}{\omega r \sin \theta} & 0 \\ 0 & 0 & \frac{z}{2\omega r} \end{vmatrix}.$$

Here and further, corresponding letters mean parameters of neutron.

Let us find rest energy of neutron.

$$\begin{aligned}
 W &= \int_V w dV = \int_V \frac{[E_r H_\theta]}{v_\varphi} dV = \\
 &= \int_0^a \int_0^\pi \int_0^{2\pi} \frac{E_0^2 \frac{a}{r} \sin^2 \varphi}{\omega r z \sin \theta} (1 + 2p \ln \frac{r}{a} + p^2 \ln^2 \frac{r}{a}) r^2 \sin \theta d\varphi d\theta dr = \\
 &= \frac{\pi^2 E_0^2 a}{\omega z} [a + 2p a \ln a - 2p \ln 0 - 2p a - 2p a \ln a + p^2 a (\ln^2 1 - 2 \ln 1 + 2) - p^2 a (0 \ln^2 0 - 2 * 0 \ln 0)] = \\
 &= \frac{\pi^2 E_0^2 a^2 (1 - 2p + 2p^2)}{\omega z}. \tag{55}
 \end{aligned}$$

Further. Charge of neutron is equal to zero.

$$q = \oint_S \varepsilon_r E_r dS = 0.$$

Really,

$$\int_0^{2\pi} \int_0^\pi \frac{E_0 \sin \varphi}{2\omega a z} a^2 \sin \theta d\varphi d\theta = 0.$$

It is obvious that

$$\int_0^\pi \int_0^\pi \frac{E_0 a \sin \varphi \sin \theta}{2\omega z} d\varphi d\theta = - \int_\pi^{2\pi} \int_0^\pi \frac{E_0 a \sin \varphi \sin \theta}{2\omega z} d\varphi d\theta \neq 0.$$

It is logical to assume that

$$\int_0^\pi \int_0^\pi \frac{E_0 a \sin \varphi \sin \theta}{2\omega z} d\varphi d\theta = - \int_\pi^{2\pi} \int_0^\pi \frac{E_0 a \sin \varphi \sin \theta}{2\omega z} d\varphi d\theta = e.$$

Then

$$\frac{2 E_0 a}{\omega z} = e. \tag{56}$$

Magnetic moment for neutron will be searched as the sum:

$$M = M_m + M_L,$$

where  $M_m$  - magnetic moment created by volume current;  $M_L$  - magnetic moment, attributed to impulse moment, i.e. to rotation.

$$M_m = \frac{1}{2} \int_0^a \int_0^\pi \int_0^{2\pi} \frac{E_0 \sqrt{a} r \sin \theta \sin \varphi}{z r^{3/2}} (p + \frac{1}{2} + \frac{1}{2} p \ln \frac{r}{a}) r^2 \sin \theta d\varphi d\theta dr = 0;$$

as

$$\int_0^{2\pi} \sin \varphi d\varphi = 0.$$

$$M_L = \gamma L,$$

$$M_L = \gamma \frac{\hbar}{2} = M. \quad (57)$$

Now we shall write down the system of equations for neutron.

$$\left. \begin{aligned} \frac{2E_0 a}{\omega z} &= e; & (56) \\ \frac{\pi^2 E_0^2 a^2}{\omega z} (1 - 2p + 2p^2) &= \hbar \omega; & (55') \\ \gamma \frac{\hbar}{2} &= -0,96623707 * 10^{-26}. & (57') \end{aligned} \right\}$$

Boundary conditions: at  $r = a$

$$\begin{aligned} \varepsilon_r &= \varepsilon_\theta = \varepsilon_0; \\ \mu_r &= \mu_\theta = \mu_0; \end{aligned}$$

hence

$$z = \sqrt{\frac{\mu_0}{\varepsilon_0}}.$$

From (54) и (54') follows that

$$\varepsilon_\varphi = \varepsilon_r \frac{2}{\sin \theta};$$

and from (54) и (54''), that

$$\mu_\varphi = \mu_\theta \frac{2}{\sin \theta}.$$

$$v_\varphi = \omega a \sin \theta = \frac{1}{\sqrt{\varepsilon_\varphi \mu_\varphi}} = \frac{1}{\sqrt{\varepsilon_0 \mu_0 \frac{4}{\sin^2 \theta}}} = \frac{c \sin \theta}{2}.$$

So

$$\omega a = \frac{c}{2};$$

$$a = \frac{c}{2\omega} = 1,0500973 * 10^{-16} (m).$$

Here  $\omega = 1,4274508 * 10^{24} \text{ Hz}$  - Compton circular frequency of neutron.

Let us solve system (56),(55'),(57').

$$E_0 = \frac{e\omega^2 \sqrt{\frac{\mu_0}{\varepsilon_0}}}{c}. \quad (56'')$$

$$E_0 = E_0^n = 4,1024444 * 10^{23} \left(\frac{V}{m}\right).$$

We substitute (56'') в (55').

$$p^2 - p + 0,5 - \frac{2\hbar}{\pi^2 e^2 \sqrt{\frac{\mu_0}{\varepsilon_0}}} = 0;$$

$$p_1 = 1,8999321;$$

$$p_2 = -0,8999321;$$

$p$  must be negative, therefore we select

$$p_2 = p_n = -0,8999321.$$

From (57') we find  $\gamma$ .

$$\gamma = -1,8324711 * 10^8 \left(\frac{1}{T * s}\right).$$

Let us write down the system of equations for antineutron.

$$\left\{ \begin{array}{l} \frac{2 E_0 a}{\omega z} = e; \\ \frac{\pi^2 E_0^2 a^2}{\omega z} (1 - 2p + 2p^2) = \hbar \omega; \\ \gamma \frac{\hbar}{2} = 0,96623707 * 10^{-26}. \end{array} \right.$$

Boundary conditions are the same, as at neutron, hence

$$z = \sqrt{\frac{\mu_0}{\varepsilon_0}};$$

$$a = \frac{c}{2\omega} = 1,0500973 * 10^{-16} (m).$$

The last system with exactness to a sign has the same solutions, as system (56),(55'),(57').

$$E_0^{\bar{n}} = E_0^n = 4,1024444 * 10^{23} \left(\frac{V}{m}\right);$$

$$p_{\bar{n}} = p_n = -0,8999321;$$

$$\gamma_{\bar{n}} = -\gamma_n = 1,8324711 * 10^8 \left(\frac{1}{T * s}\right).$$

## Conclusion

Within the framework of the model, which is considered, electron, proton and neutron represent a monochromatic electromagnetic wave of corresponding frequency spread along parallels inside the spherical area, i.e. a wave, rotating around some axis. At that along each parallel, exactly half of wave length for electron and proton and exactly one wave length for neutron, is kept within, thus this is rotating soliton. This is caused by presence of dispersion and anisotropy of a strictly defined type inside the particles. In electron vector  $E$  is directed to centre of particle, that correspond to negative charge, and in proton vector  $E$  is directed from centre of particle, that correspond to positive charge. Thus, by natural way, all basic parameters of particles are obtained: charge, rest energy, mass, radius, magnetic moment and spin, that is confirmed by mathematical expressions, which are discovered.

## Literature

1. L. I. Sarycheva. Structure of matter. Sorosovskiy obrazovatelniy jurnal. Volume 6, № 2, 2000. WWW.issep.rssi.ru.
2. D. I. Blokhintsev. Principles of quantum mechanics. - M.: Nauka, 1983.
3. M. Born. Atomic physics. - M.: Mir, 1967.
4. M. M. Bredov, V. V. Rumyantsev, I. N. Toptygin. Classical electrodynamics. Edited by I. N. Toptygin.: - M.: Nauka, 1985.
5. M. B. Vinogradova, O. V. Rudenko, A. P. Sukhorukov. Theory of waves. - M.: Nauka, 1990.
6. A. D. Vlasov, B. P. Murin. Physical quantities units in science and technology. Reference manual. - M.: Energoatomizdat, 1990.
7. S. K. Godunov. Mathematical physics equations. - M.: Nauka, 1979.
8. S. G. Kalashnikov. Electricity. - M.: Nauka, 1985.
9. E. Kamke. Reference manual on ordinary differential equations. Translated from German by S.V. Fomina. - M.: Nauka, 1976.
10. L. D. Landau, E. M. Lifshits. Quantum mechanics. Non-relativistic theory. - M.: Nauka, 1989.

11. L. D. Landau, E. M. Lifshits. Field theory. - M.: Nauka, 1973.
12. L. D. Landau, E. M. Lifshits. Electrodynamics of continuous mediums. - M.: Nauka, 1982.
13. A. P. Prudnikov, Yu. A. Brychkov, O. I. Marychev. Integrals and series. Elementary functions. - M.: Nauka, 1981.
14. B. Taylor, V. Parker, D. Langerberg. Fundamental constants and quantum electrodynamics. Translated from English by Candidates of physical and mathematical sciences V.D. Burlakov, V.G. Krechet and V.G. Lapchinskiy. Edited by Professor B. A. Mamyrin - M.: Atomizdat, 1972.
15. Ya. P. Terletskiy, Yu. P. Rybakov. Electrodynamics. - M.: Vysshaya Shkola, 1990.
16. E. V. Shpol'skiy. Atomic physics. Volume 1. Introduction in atomic physics. - M.: Nauka, 1984.
17. E. V. Shpol'skiy. Atomic physics. Volume 2. Principles of quantum mechanics and structure of electronic cover of atom. - M.: Nauka, 1984.
18. A. G. Kyriakos. The electrodynamics form concurrent to the Dirac electron theory. Physics Essays, volume 16, number 3, 2003.
19. A. G. Kyriakos. The massive neutrino-like particle of the non-linear electromagnetic field theory. Apeiron, Vol. 12, No 1, January 2005.
20. A. G. Kyriakos. Yang-Mills equation as the equation of the superposition of the non-linear electromagnetic waves. <http://arXiv.org/abs/hep-th/0407074>, 09.07.2004.
21. Alexander G. Kyriakos. Non-linear Theory of quantized Electromagnetic Field equivalent to the Quantum Field Theory. <http://www.partphys.envy.nu>.
22. I. P. Ivanov. Quark model is not quite correctly?! <http://www.astronet.ru/db/msg/1188217> , 25.03.2003, in Russian.
23. V. Kopelyowicz. Topologic soliton models of baryons and its predictions. Scientific.ru, in Russian.
24. D. Diakonov, V. Petrov and M. Polyakov, Z. Phys. A359, 305 (1997).
25. T. H. R. Skyrme, Nucl. Phys. 31 (1962), 556.
26. M. Polyakov et al., Eur. Phys. J. A9, 115 (2000).
27. LEPS Collaboration: T. Nakano, et al., Evidence for Narrow  $S = +1$ . Baryon Resonance in Photo-production from Neutron, Phys. Rev. Lett. 91 (2003) 012002.

Hep-ex/0301020.

28. M. V. Polyakov, A. Rathke, On photoexcitation of baryon antidecuplet, Eur. Phys. J. A 18 (2003), 691-695, hep-ph/0303138, 17 March 2003.

29. D. Diakonov and V. Petrov. Baryons as solitons. Elementary particles. Moscow. Energoatomizdat, 1985, vol. 2, p. 50, in Russian.

30. M. Chemtob, Nucl. Phys. B256 (1985) 600.

31. M. Praszalowicz, in Skyrmons and Anomalies, M. Jezabek and M. Praszalowicz, eds., World Scientific (1987) p. 112.

32. H. Walliser, in Baryon as Skyrme Soliton, p. 247, ed. by G. Holzwarth, World Scientific, 1992;

H. Walliser, Nucl. Phys. A548 (1992) 649.

33. D. Diakonov, V. Petrov and M. Polyakov, Exotic Anti-Decuplet of Baryons: Prediction from Chiral Solitons, Z. Phys. A359 (1997), 305-314, arXiv: hep-ph/9703373.

34. I. P. Ivanov. Last days of Standard Model?

<http://www.scientific.ru/journal/news/0702/n140702>, 14.07.2002.

P. S. Further researches on the basis of results, which were obtained, intend solution of following tasks:

1. Elaboration of physic-mathematical model of photon and neutrino structure.
2. Elaboration of physic-mathematical model of atomic nuclei structure for all chemical elements.

It is my firm belief that solution of this tasks will assist to achieve great leap in such fields, as nuclear power engineering, nanotechnology and high-powerful lasers.